

HOW FAINT CAN WE GO?

mag 14 ... 15 ... 16 ... 17 ? ... 18 ??

**Exploring the magnitude limits
of amateur spectroscopy
using a modified ALPY spectrograph
at Three Hills Observatory**

HOW FAINT CAN WE GO?

Modifying the ALPY Spectrograph for fainter objects

Robin Leadbeater



THREE HILLS OBSERVATORY UK

Celestron C11 on EQ6 mount
(EQMod, ASCOM, Cartes du Ciel)

Spectrographs:-

LHIRES III (150,600,1200,2400)

Star Analyser (100, 200)

ALPY 600

Remotely operated via wireless network
(Windows Remote Desktop)

ALPY SPECTROGRAPH

Focal Reducer
f6

Guide Camera
ATIK 16 IC-S
(cooled ICX415AL)


Science Camera
ATIK 314L+
(cooled ICX285AL)

Calibration Module
Tungsten &
Ne/Ar/H lamps


USB
remote control for
calibration module

Guiding Module
23um mirror slit

Core Module
Collimator
Grism
Camera lens



MODIFYING THE ALPY SPECTROGRAPH (ALPY 200)



Removing the the Grism from
the ALPY Core Module

Taking care that the collimator
lens does not fall out!




Standard
600 l/mm
Grism



Modified
200 l/mm Grism
(Made using the
Star Analyser 200
master grating)

BENCH TESTS - ALPY 200 v 600


Resolution



The dispersion and resolution of the ALPY200 are ~ 4 lower compared with the ALPY 600

R = 130 at 6000 A is sufficient for supernova identification for example

Efficiency



The efficiency of the 200 l/mm grism is ~ 2x that of the 600 l/mm grism

Combining this with the 4x reduction in dispersion results in an **8x brighter spectrum**

ADVANTAGES OF THE ALPY 200

Compared with the Star Analyser

The slit reduces the sky background level by a factor of ~ 100 , reducing sky noise and eliminating interference from field stars and their spectra.

The mirror slit guider allows longer individual exposures, reducing the total camera read noise contribution.

The fully collimated optics produce a sharp spectrum over the full wavelength range, allowing a lower dispersion to be used for the same resolution.


Compared with the ALPY 600

The higher efficiency of the 200 l/mm grism increases the flux in the spectrum.

The reduced dispersion minimises the camera thermal noise contribution compared with increasing the effective pixel size by in camera binning


FIRST RESULTS -

QSO 3C273 (mag 12.5) 20 deg from full moon




Raw spectrum with H beta emission line visible against the bright lunar spectrum and Na D from street lights




Reduced spectrum showing red shifted Balmer lines ($z = 0.158$)




Comparison with professional spectrum



QSO APM 08279 +5255 (mag 15.5 and 12GLyr)



The key features in the spectrum are clear even in a single 600s exposure. The features at ~6000Å and ~7500Å are red shifted from the UV ($z=3.91$, 12 GLyr). The absorption at ~5000Å is from a gravitationally lensing object in the line of sight at $z=3.07$




Comparison with professional spectrum

Light pollution spectrum at THO compared with VLT Paranal
Note the natural airglow molecular bands in IR from OH


SN 2014AS in NCG 5410 (type 1c mag 16.5)

Discovered by amateur Dave Grennan 2014- 04-18



Guide camera image.


The sn was not visible during guiding so a mag 13 field star was used



Spectrum image before and after sky subtraction

Spectra of the supernova and galaxy core above it. Note the clear H alpha emission from the galaxy core and OH air glow bands in the IR

THREE HILLS OBSERVATORY ALPY 200 2014-04-26
Black - PSN J14005449+4058596
Blue - NGC 5410 measured $z=0.0129$ (H alpha)




Reduced spectra for galaxy and supernova.

Exposure 110 min SNR ~30

The sn spectrum does not show strong H alpha emission or Si 6150 absorption features (not type II or 1a)


The measured red shift of NGC5410 (Ha) = 0.0124 (published figure = 0.0129)





GELATO

supernova identification program suggests a best fit to type 1b/c


PSNJ15024996+4847062_20140521 in PGC 2325560 (mag ~16)
Discovered by amateurs R. Gagliano, J. Newton T. Puckett 2014- 05-13



Discovery image




Spectrum image before and after sky subtraction
 Spectrum of the supernova (centre).
 The lower spectrum is a field star caught in the slit



Reduced spectrum (red)

overlaid on professional
 spectrum (blue) taken 6
 days earlier (ATel 6140)

The spectrum shows a
 strong Si II 6150 absorption
 feature redshifted to 6300 Å
 (characteristic of type Ia)



— @psn15024996+4847062_wlcal_respcor.fit z:0.026 (V_{orig})
 — 2009dc type:Ia pec phase:7.2d rel.to Bmax obs.date:20090502 z:0.02163 (flux scaled)

GELATO

supernova identification program
 gives an excellent fit to type Ia
 a few days after maximum